

# Ephemeris Updates for Seven Selected HATNet Survey Transiting Exoplanets

A. Poro<sup>1,\*</sup>, F. Ahangarani Farahani<sup>2</sup>, E. Jahangiri<sup>2</sup>, A. Sarostad<sup>2</sup>, M. Gozarandi<sup>2</sup>, M. Haghgou<sup>2,3</sup>, F. Abolhassani<sup>2</sup>, A. Fakhrabadi<sup>2</sup>, Y. Jongen<sup>4</sup>, A. Wünsche<sup>5</sup>, R. Naves<sup>6</sup>, P. Guerra<sup>7</sup>, A. Marchini<sup>8</sup>, M. Salisbury<sup>9</sup>, R. Ehrenberger<sup>10</sup>, V-P. Hentunen<sup>11</sup>

<sup>1</sup>Astronomy Department of the Raderon AI Lab., BC., Burnaby, Canada

<sup>2</sup>BSN Project, Transiting Exoplanets Department, Tehran, Iran

<sup>3</sup>Department of Chemistry, Faculty of Science, University of Guilan, Rasht, Iran

<sup>4</sup>Observatoire de Vaison-La-Romaine, Departementale 51, 84110 Vaison-La-Romaine, France

<sup>5</sup>Observatoire des Baronnies Provençales, Route de Nyons, F-05150 Moydans, France

<sup>6</sup>Observatorio Montcabrer, Cabrils, Spain

<sup>7</sup>Observatori Astronòmic Albanyà, Camí de Bassegoda S/N, Albanyà E-17733, Girona, Spain

<sup>8</sup>University of Siena - Astronomical Observatory, Via Roma 56, 53100 Siena, Italy

<sup>9</sup>British Astronomical Association, 25 Farringdon Street, London EC4A 4AB, England

<sup>10</sup>Czech Astronomical Society, Fričova 298, 251 65 Ondřejov, Czech Republic

<sup>11</sup>Taurus Hill Observatory, 79480 Varkaus, Finland

Accepted —. Received —; in original form —

## ABSTRACT

We refined the ephemeris of seven transiting exoplanets HAT-P-6b, HAT-P-12b, HAT-P-18b, HAT-P-22b, HAT-P-32b, HAT-P-33b, and HAT-P-52b. We observed 11 transits from eight observatories in different filters for HAT-P-6b and HAT-P-32b. Also, the Exoplanet Transit Database (ETD) observations for each of the seven exoplanets were analyzed, and the light curves of five systems were studied using Transiting light Exoplanet Survey Satellite (TESS) data. We used Exofast-v1 to simulate these ground- and space-based light curves and estimate mid-transit times. We obtained a total of 11, 175 and 67 mid-transit times for these seven exoplanets from our observations, ETD and TESS data, respectively, along with 155 mid-transit times from the literature. Then, we generated transit timing variation (TTV) diagrams for each using derived mid-transit times as well as those found in the literature. The systems' linear ephemeris was then refined and improved using the Markov Chain Monte Carlo (MCMC) method. All of the studied exoplanets, with the exception of the HAT-P-12b system, displayed an increasing trend in the orbital period in the TTV diagrams.

**Key words:** planetary systems – planets and satellites: gaseous

## 1 INTRODUCTION

The number of exoplanets discovered and characterized each year has been increasing since the results of the first exoplanet detection [1]. Hot Jupiters are an important type of planetary gas giant with masses and radii similar to Jupiter but orbiting their host stars with short orbital periods (most less than 10 days), making them a good target system to discover and study [2][3]. The transit technique is the most efficient way to improve our understanding of exoplanets through ground- and space-based surveys. Furthermore, photometric transit surveys combined with radial velocity data have become one of the most successful methods for detecting transiting exoplanets over the past decade [4]. High-precision transit observations provide information to refine planetary parameters such as the planet's size, mass, atmosphere, and orbital ephemerides [5][6]. Moreover, photometric

transit surveys allow us to study the variations of the orbital periods through TTV analysis. Space telescopes have longer available observational time, and they are not affected by the Earth's atmosphere as well. TESS is one of the most significant space-based survey missions for the discovery and observation of transiting exoplanets. TESS was launched in 2018 to observe new exoplanets orbiting bright nearby stars that are brighter than Kepler mission stars [7]. Furthermore, when combined with previous work, this space mission provides precise transit timing for discovered exoplanets, which is critical for obtaining a better transit ephemeris [8].

Based on our observations, TESS, ETD<sup>1</sup>, and literature observations, we updated orbital ephemeris for the HAT-P-6b, HAT-P-12b (TESS ID 198108326), HAT-P-18b (TESS ID 21744120), HAT-P-22b (TESS ID 252479260), HAT-P-32b, HAT-P-33b (TESS ID

\* Corresponding author: poroatila@gmail.com

<sup>1</sup> <http://var2.astro.cz/ETD/>

239154970), and HAT-P-52b (TESS ID 436875934). These exoplanets were discovered by the Hungarian-made Automated Telescope Network (HATNet) survey.

## 2 OBSERVATIONS AND METHOD

### 2.1 Observation and data reduction

Observations in this study have been made regarding exoplanets HAT-P-6b and HAT-P-32b during the years 2018 to 2022. A total of nine observation nights have been done for these two exoplanets; five and four nights for HAT-P-6b and HAT-P-32b, respectively. All these photometric observations have been done with small telescopes at eight observatories. We used CCD and standard filters in these observations. The information about the observatories, telescopes, CCDs, and data reduction software that we used is listed in Table 1. In Table 1, an abbreviated name has been determined for each observatory just to identify them in this study. The basic data reduction for the dark, bias, and flat field of each CCD image was carried out in accordance with the standard technique.

### 2.2 ETD data

To obtain the refined orbital ephemeris of selected HATNet exoplanets, we also collected also light curves, which were sourced from astronomers through the ETD archive [9]. Light curves were obtained from various filters and time scales. We used data in ETD that we were confident enough to be appropriate; for example, we did not use data whose declared time was less than three digits. We used those which generally have a quality index (DQ) of less than three [9]. All times in the data were converted from JD or HJD to  $BJD_{TDB}$  based on the geographic location of observation and RA(J2000) and DEC(J2000) from the Simbad<sup>2</sup> astronomical database.

In some ETD light curves, the airmass effect has been ignored, so airmass must be calculated based on the observers' location, which influences and improves the measured mid-transit times of related light curves. Therefore, we computed the airmass using the Astropy package in Python [10].

### 2.3 TESS data

Five of these exoplanets were observed by TESS, and HAT-P-6b, and HAT-P-32b have no TESS data yet. TESS observed the five host stars at 120-second cadences. We collected TESS data from the Mikulski space telescope archive (MAST). TESS style curves were extracted by LightKurve<sup>3</sup> code from the MAST Python package.

### 2.4 Method

We relied on the AstroImageJ software [11] to normalize all of the data. Figure 1 shows the folded TESS light curves for five selected exoplanets. Finally, all ground- and space-based light curves were applied to Exofast-v1 [12] for modeling purposes; as a consequence, the output mid-transit times and associated uncertainties were employed. Figure 2 provides an example of a modeled observation of TESS and this study's observation. The extracted transit mid-times from our observations and TESS data are provided in Tables

2 and 3. Tables 5-11 include the literature and ETD transit mid-times.

We plotted TTV diagrams for seven selected exoplanets using derived mid-transit times and those available within the literature. Our MCMC analysis of these timings enabled us to refine the linear ephemeris of the systems. We applied the MCMC method, or sampling from the posterior probability distributions of the coefficients (100 walkers, 10000 step number, and 1000 burn-in) using the Pymc3 package in Python [13]. Figure 3 shows all TTV-diagrams of studied exoplanets and also displays the posterior distributions for the fitted parameters using the MCMC method (dT and dP).

## 3 REFINED TRANSIT EPHEMERIS

Follow-up observations of known transiting exoplanets, either with photometry or high-resolution spectroscopy, are an important issue. Accordingly, a growing quantity of this data provides the refined physical planetary properties, formation, and evolution processes of these exoplanets. Since the orbital ephemeris of exoplanets demonstrates their places in their orbits, refinement of the exoplanet's ephemeris would be a key factor for the prediction of future transit events.

Based on the reference ephemeris for each exoplanet, epochs, and O-Cs are calculated. According to extracted mid-transit times, we can compute a new ephemeris by a well-known linear relation (1),

$$T_c = T_0 + E \times P \quad (1)$$

where  $T_0$  is considered as the reference mid-transit time,  $P$  and  $E$  are the orbital period and the number of epochs passed since  $T_0$ , respectively. New ephemeris and reference ephemeris for exoplanets are given in Table 4.

### 3.1 HAT-P-6b

The exoplanet HAT-P-6b is a hot Jupiter with a radius of  $1.33 \pm 0.06 R_J$  and a mass of  $1.06 \pm 0.12 M_J$  [14]. The [14] study, which was based on the primary transit method's results, announced its discovery. According to a report by [14] on the characteristics of the planet and its host star,  $e=0.046 \pm 0.031$  indicates that the planet's orbit is almost circular. [19] collected transit light curves for this exoplanet and refitted available photometry data with their new and literature mid-transit times. A revised orbital period and mid-transit time for this exoplanet were also updated by [20].

We collected the high-quality mid-transit times that were previously published in the literature, and the mid-transit times resulted from the modeling of ETD light curves for plotting an updated TTV diagram. We extracted a total of six mid-transit times from our observations and 13 mid-transit from ETD. We note that this exoplanet has not had TESS data.

### 3.2 HAT-P-12b

[15] used the transit method to discover the exoplanet HAT-P-12b, which orbits a dwarf star. According to the results of the discovery paper, HAT-P-12b is one of the least massive gas giant planets that was found until its discovery and is an exoplanet of the sub-Saturn type. The  $J$  band photometry study by [21] of this exoplanet obtained the precise mid-transit time to constrain the transit-time variations of the HAT-P-12b system. [22] observed three new transit light

<sup>2</sup> <http://simbad.u-strasbg.fr/simbad/>

<sup>3</sup> <https://docs.lightkurve.org/>

**Table 1.** The observatories of this study and the instruments that were employed.

Observatory	Telescope	CCD	Data reduction Software
Rasteau Observatory, France (RO)	PlaneWave CDK 17"	SBIG STXL11004	Muniwin / C-munipack
Montcabrer private observatory, Spain (MO)	SCT 12"	SBIG ST8-XME	Fotodif
Observatori Astronomic Albanyà, Spain (AA)	Meade ACF 16"	Moravian Instruments G4-9000	Fotodif
Astronomical Observatory, University of Siena (K54), Italy (AO)	MCT 300 mm	SBIG STL-6303	Muniwin / C-munipack
Observatoire des Baronnies Provençales, France (BO)	Cassegrain 430 mm	Zwo ASI6200 Pro mono	Muniwin / C-munipack
Private Observatory, Czech Republic (PO)	400 mm	SBIG ST-10 XME	AstroImageJ 3.2.10
Crow-Observatory Vranová, Czech Republic (CO)	NWT 300 mm	Moravian Instruments G2-3200	Muniwin / C-munipack
Taurus Hill Observatory, Finland (TO)	SCT 14"	SBIG ST-8 XME	AIP4Win v2.4.10

**Table 2.** Extracted ground-based transit times for HAT-P-6b and HAT-P-32b in this study.

Exoplanet	Observatory	$T_c(BJD_{TDB})$	Error	Filter	Eopch	O-C
HAT-P-6b	MO	2455430.46458	0.00110	Optec $R$ cousins	362	0.0075
HAT-P-6b	RO	2458312.50536	0.00063	Baader imaging $G$	1110	0.0155
HAT-P-6b	AA	2458389.56389	0.00101	Baader $J - CV$	1130	0.0143
HAT-P-6b	AO	2459441.43053	0.00101	Johnsons-Cousins $I$	1403	0.0161
HAT-P-6b	RO	2459441.43128	0.00095	Baader imaging $R$	1403	0.0168
HAT-P-6b	RO	2459468.40283	0.00124	Baader imaging $R$	1410	0.0175
HAT-P-32b	CO	2459107.46235	0.00097	Johnsons-Cousins $R_c$	2180	-0.0015
HAT-P-32b	TO	2459191.31427	0.00034	Baader Bessel photometric $R$	2219	0.0001
HAT-P-32b	BO	2459507.36578	0.00017	Johnsons-Cousins $V$	2366	0.0005
HAT-P-32b	PO	2459593.36707	0.00024	Johnsons-Cousins $R_c$	2406	0.0015
HAT-P-32b	BO	2459593.37024	0.00041	Johnsons-Cousins $R_c$	2406	0.0046

curves of HAT-P-12b, and with existing literature data, they came up with an improved ephemeris. [22] also refined the absolute physical properties of the star-planet system. [23] updated the ephemeris of HAT-P-12b according to six transits for this system by applying a least-squares linear fit to all available transit times. According to their results, no long-term TTVs were apparent. [24] studied HAT-P-12b in bands  $V$  and  $I$  to investigate the transmission spectrum of this system. [24] observed 23 new photometric transit light curves, and analysis showed no indication of star-spot influence on the calculated transit parameters. [25] studied this exoplanet's atmosphere. In fact, the goal of this research was to specify an appropriate solution for future studies of other exoplanetary atmospheres. Spectroscopic observations using the Large Binocular Telescope (LBT) were done by [26] to obtain an atmosphere transmission spectrum of this exoplanet. They found no evidence of Na or K absorption features in the relatively flat transmission spectrum, which is in agreement with the HST transmission spectrum. Furthermore, [27] included six new mid-transit times to determine a new ephemeris by a linear fit to a satisfactory level. [28] also reported an infrared transmission photometry of HAT-P-12b with the other 48 exoplanets with the largest analysis of Spitzer/IRAC observations to study the influence of infrared photometry on atmospheric chemical properties.

We used mid-transit times conducted from the modeling of ETD light curves and TESS in association with data published in previous literature for plotting a new TTV diagram. We extracted 27 mid-transit times from ETD and 6 mid-transit times from sector 23 of TESS for HAT-P-12b.

### 3.3 HAT-P-18b

HAT-P-18b is a low-density Saturn-mass exoplanet orbiting a super-solar metallicity K2 dwarf star [29]. The discovery observations of this exoplanet have been made by [29] using the transit method to obtain the orbital and physical properties of the system. [29] reported a non-zero ( $e = 0.0840.048$ ) eccentricity for HAT-P-18b. Complementary new photometric observations of the full transit were also analyzed by [30] in order to independently estimate the parameters of the host star and HAT-P-18b. [31] performed the TTV study of HAT-P-18b with a limited number of existing high-quality data and they presented ground-based transmission spectroscopy of HAT-P-18b. This exoplanet was described as a hot Jupiter by [32], who also found Rayleigh-scattering in the atmosphere, and their results confirmed that ground-based observations are suitable to determine the opacity sources of exoplanets' atmospheres. [33] observed HAT-P-18b as a part of the original research by the young twinkle students (ORBYTS) program to refine its transit ephemerides. The atmosphere of this exoplanet has been studied by [34]. Moreover, [35] derived the refined ephemeris from observations provided by the ExoClock network in combination with previous literature data.

For HAT-P-18b, we obtained seven mid-transit times from sectors 25 and 26 of TESS, and 21 mid-transit times from ETD.

### 3.4 HAT-P-22b

[17] reported the discovery of the exoplanet HAT-P-22b. It is among the moderately massive and compact hot Jupiters, orbiting a fairly metal-rich dwarf star with a  $V=9.732$  magnitude. [36] presented the first photometric follow-up observation of bright transiting exoplanets by using a defocusing technique. Following this, [37] performed a

**Table 3.** Extracted TESS transit times for five exoplanets.


Exoplanet	$T_c(BJD_{TDB})$	Error	Epoch	O-C	Exoplanet	$T_c(BJD_{TDB})$	Error	Epoch	O-C
HAT-P-12b	2458933.54320	0.00053	1405	-0.0014	HAT-P-33b	2459506.13961	0.00051	1265	0.0040
HAT-P-12b	2458936.75601	0.00052	1406	-0.0016	HAT-P-33b	2459509.61506	0.00052	1266	0.0050
HAT-P-12b	2458939.96969	0.00048	1407	-0.0010	HAT-P-33b	2459516.56424	0.00058	1268	0.0053
HAT-P-12b	2458946.39512	0.00055	1409	-0.0017	HAT-P-33b	2459520.03824	0.00054	1269	0.0048
HAT-P-12b	2458949.60784	0.00050	1410	-0.0020	HAT-P-33b	2459523.51230	0.00048	1270	0.0044
HAT-P-12b	2458952.82114	0.00048	1411	-0.0018	HAT-P-33b	2459526.98761	0.00052	1271	0.0052
HAT-P-18b	2458989.25305	0.00059	776	0.0055	HAT-P-33b	2459530.46206	0.00057	1272	0.0052
HAT-P-18b	2458994.76038	0.00048	777	0.0048	HAT-P-33b	2459533.93584	0.00055	1273	0.0045
HAT-P-18b	2459005.77787	0.00050	779	0.0062	HAT-P-33b	2459537.41024	0.00057	1274	0.0044
HAT-P-18b	2459011.28554	0.00051	780	0.0059	HAT-P-33b	2459540.88529	0.00065	1275	0.0050
HAT-P-18b	2459016.79255	0.00059	781	0.0048	HAT-P-33b	2459544.35953	0.00062	1276	0.0048
HAT-P-18b	2459027.80975	0.00060	783	0.0060	HAT-P-33b	2459547.83335	0.00063	1277	0.0041
HAT-P-18b	2459033.31868	0.00060	784	0.0069	HAT-P-33b	2459554.78399	0.00053	1279	0.0058
HAT-P-22b	2458871.63055	0.00020	1227	0.0166	HAT-P-33b	2459558.25712	0.00055	1280	0.0044
HAT-P-22b	2458874.84229	0.00018	1228	0.0161	HAT-P-33b	2459561.73071	0.00057	1281	0.0036
HAT-P-22b	2458878.05478	0.00018	1229	0.0164	HAT-P-33b	2459568.68054	0.00053	1283	0.0044
HAT-P-22b	2458881.26687	0.00019	1230	0.0163	HAT-P-33b	2459572.15466	0.00054	1284	0.0041
HAT-P-22b	2458887.69120	0.00017	1232	0.0161	HAT-P-33b	2459575.63036	0.00053	1285	0.0053
HAT-P-22b	2458890.90361	0.00020	1233	0.0163	HAT-P-33b	2459582.57829	0.00053	1287	0.0043
HAT-P-22b	2458894.11620	0.00018	1234	0.0167	HAT-P-33b	2459586.05330	0.00053	1288	0.0048
HAT-P-22b	2458897.32739	0.00020	1235	0.0157	HAT-P-33b	2459589.52750	0.00051	1289	0.0046
HAT-P-22b	2459613.65575	0.00019	1458	0.0190	HAT-P-33b	2459599.95038	0.00055	1292	0.0040
HAT-P-22b	2459616.86844	0.00018	1459	0.0194	HAT-P-33b	2459603.42543	0.00053	1293	0.0046
HAT-P-22b	2459620.08042	0.00018	1460	0.0192	HAT-P-52b	2459475.83726	0.00205	1316	0.0026
HAT-P-22b	2459626.50524	0.00016	1462	0.0196	HAT-P-52b	2459478.59303	0.00204	1317	0.0048
HAT-P-22b	2459629.71736	0.00016	1463	0.0195	HAT-P-52b	2459489.60804	0.00179	1321	0.0054
HAT-P-22b	2459632.92936	0.00017	1464	0.0193	HAT-P-52b	2459492.35880	0.00239	1322	0.0026
HAT-P-33b	2458845.98935	0.00055	1075	0.0038	HAT-P-52b	2459495.11585	0.00202	1323	0.0060
HAT-P-33b	2458849.46501	0.00051	1076	0.0050	HAT-P-52b	2459500.61943	0.00171	1325	0.0024
HAT-P-33b	2458852.93919	0.00053	1077	0.0047	HAT-P-52b	2459503.37368	0.00163	1326	0.0031
HAT-P-33b	2458859.88850	0.00051	1079	0.0051	HAT-P-52b	2459506.12841	0.00149	1327	0.0042
HAT-P-33b	2458863.36179	0.00051	1080	0.0039	HAT-P-52b	2459508.87918	0.00163	1328	0.0014
HAT-P-33b	2458866.83641	0.00054	1081	0.0041	HAT-P-52b	2459514.38955	0.00156	1330	0.0045
HAT-P-33b	2459502.66693	0.00052	1264	0.0058					

**Table 4.** The new ephemeris derived by a linear fit on the TTV diagram of each exoplanet and reference ephemeris for computing epochs and the TTV values.


Exoplanet	New ephemeris ( $BJD_{TDB}$ )	Reference ephemeris ( $BJD_{TDB}$ )
HAT-P-6b	$2454035.6769526(3) + 3.85300(15) \times E$	$2454035.67652(28) + 3.852985(5) \times E$ [14]
HAT-P-12b	$2454419.19585(6) + 3.21305852(8) \times E$	$2454419.19556(20) + 3.2130598(21) \times E$ [15]
HAT-P-18b	$2454715.022802(97) + 5.5080288(2) \times E$	$2454715.02174(20) + 5.508023(6) \times E$ [16]
HAT-P-22b	$2454930.22043(16) + 3.2122330(1) \times E$	$2454930.22001(25) + 3.212220(9) \times E$ [17]
HAT-P-32b	$2454420.44713(6) + 2.15000821(5) \times E$	$2454420.44637(9) + 2.150008(1) \times E$ [16]
HAT-P-33b	$2455110.92683(12) + 3.4744769(1) \times E$	$2455110.92595(22) + 3.474474(1) \times E$ [16]
HAT-P-52b	$2455852.10370(23) + 2.7535989(3) \times E$	$2455852.10326(41) + 2.7535953(94) \times E$ [18]

follow-up transit observation using a defocusing technique and they derived one complete transit and computed the mid-transit times for HAT-P-22b. The near-UV and optical photometric observations of HAT-P-22b were made by [38] to study the atmosphere of this exoplanet. [38] also refined the planetary parameters and ephemerides of HAT-P-22b hot Jupiter. Accordingly, all derived parameters were in agreement with the discovery values by [17], and any non-spherical asymmetries were not seen in their data.

In order to plot a TTV diagram for this exoplanet, we extracted 30 and 14 mid-transit times resulting from the modeling of ETD and TESS light curves (sectors 21 and 48), respectively, as well as data from previous publications.



**Figure 1.** Folded TESS light curves in each sector of all selected exoplanets were obtained from the LightKurve code.




**Figure 2.** Left: HAT-P-33b observational and theoretical light curves using TESS sector 45 data; Right: The observational light curve of HAT-P-32b from this study in the V filter and the theoretical light curve.

### 3.5 HAT-P-32b

The planet HAT-P-32b was discovered by the HATNet survey in 2011 and it is a hot Jupiter exoplanet orbiting a late-F-early-G dwarf star with  $V=11.289$  magnitude. In this discovery, radial velocity measurements were taken with High-Resolution Echelle Spectrometer

and [39] transit model was used in order to describe the HATNet photometry [16]. [21] presented a JH-band photometry observation of HAT-P-32b and extracted precise mid-transit times. [21] declared that HAT-P-32b system parameters were in agreement with those reported in the [16] study and derive a period of this exoplanet with im-



**Figure 3.** The TTV diagrams of seven studied exoplanets with the linear fit on the data points and posterior distributions for the fitted parameters using MCMC ( $dT$  and  $dP$ ).

proved uncertainty. Following this, [40] reported two primary transits of HAT-P-32b during Gemini-North Gemini Multi-Object Spectrograph observations. They used white light curve analysis in order to refine the parameters of this exoplanet and derive new ephemeris. [41] updated the system properties by analyzing the results of 45 transit observations, which were observed by using the young exoplanet transit initiative (YETI) network. Moreover, [41] studied the TTV diagram to investigate the existence of an additional planet in the HT-P-32b system. [42] performed a global fit for the HAT-P-32b system based on their new photometric observations and previously published RV data in order to update the system parameters. [42] also analyzed the TTV diagram for this system and according to the results, there was no significant TTV signals. Some follow-up high-quality observations of this exoplanet were done with small observatories operated by citizen scientists in 2020 [43].

The accurate mid-transit times for HAT-P-32b were obtained from

the available data for plotting an updated TTV diagram. We extracted a total of five and 72 mid-transit times from our observations and the ETD, respectively.

### 3.6 HAT-P-33b

The planet HAT-P-33b was among the first exoplanets discovered by the HATNet survey in 2011 and was confirmed by high-precision photometry and additional radial velocity measurements [16]. HAT-P-33b is an inflated hot Jupiter orbiting a Late-F dwarf star with a short orbital period. [16] reported that HAT-P-33b has a radius of  $\sim 1.7 R_J$  which is among one of the largest measured radius for all transiting exoplanets. HAT-P-33b also has an equilibrium temperature of more than 1600 K, which is the result of the high luminosity of its host star. The TTV study of HAT-P-33b was analyzed by the transiting exoplanet monitoring project (TEMP) in the study of [44].

[44] refined HAT-P-33b orbital ephemerides and the system parameters through R-band photometric observations. Furthermore, the extended radial velocity measurements of this planet have been made by [45]. Their obtained transit parameters were consistent with the resulting parameters from [16], but smaller values for the ingress/egress duration, and the planet to star radius ratio ( $R_P/R_*$ ) were deduced. These differences were because of derived complete light curves, which led to shorter and shallower transit shapes. However, they found no sign of anomalies in the TTV diagram. [46] performed additional follow-up ground-based photometric observations to confirm the HAT-P-33b planetary parameters through TEP modeling methods. [46] used a weighted linear least-squares analysis to update the reference ephemeris for this planet. Moreover, they found that the *R*-band transit depth in HAT-P-33b is larger than its discovery value [16] when discussing the variation in planetary radius against wavelength. [46]'s photometric data revealed no evidence of star-spot for this exoplanet.

We used the mid-transit times collected from the modeling of ETD and TESS light curves to prepare a new TTV diagram for HAT-P-33b. For this exoplanet, 30 mid-transit times from TESS sectors 20, 44, 45, 46, and 47 were extracted, also with seven mid-transit times from ETD observations.

### 3.7 HAT-P-52b

HAT-P-52b is a short-period gas-giant hot Jupiter discovered and characterized in 2015 by the transit method [18]. [47] reported new photometric light curves for HAT-P-52b and refined this system's parameters by analyzing their light curves, previous photometric, and Doppler velocimetric data. [47] also performed the TTV study for HAT-P-52b and their results represented no significant trend in the TTV diagram and the measured mid-transit time was consistent with their updated linear ephemerides.

We plotted the updated TTV diagram based on the seven mid-transit times from the modeling of ETD and 10 mid-transit times from the TESS light curves (sectors 43 and 44), along with published data in previous literature.

## 4 SUMMARY AND CONCLUSION

We conducted a study on seven HATNet survey-selected transiting exoplanets and plotted the TTV diagrams. The goal of this study is to improve the planetary systems' reference ephemerides and to discuss the reasons for the period variations in these systems for future studies.

We have presented a new ephemeris for each of the seven exoplanets. For this purpose, we utilised the mid-transit times found in the literature as well as the light curves observed by ETD, TESS, and our ground-based observations. We used 11 mid-transit times from our observations in this study, which were made at eight observatories from 2018 to 2022.

Exofast-v1 was used to model the available light curves and extract the mid-transit times. We used the MCMC method to plot new TTV diagrams and refine exoplanets' ephemeris. The TTV diagrams show the orbital periods of exoplanets HAT-P-6b, HAT-P-18b, HAT-P-22b, HAT-P-32b, HAT-P-33b, and HAT-P-52b are increasing, whereas exoplanet HAT-P-12b has a declining tendency. It is probable that the six exoplanets' orbital periods increased since their ephemeris accuracy has become inaccurate over time.

According to the new ephemeris for exoplanet HAT-P-6b, it seems

that the uncertainties of  $t$  and  $P$  should be more carefully considered in future investigations and observations.

## ACKNOWLEDGEMENTS

We made use of the ETD data for this investigation. ETD is maintained by the Czech Astronomical Society (CAS), an old astronomical organization in the Czech Republic with more than hundreds of members (amateur and professional astronomers included).

This work includes data from the TESS mission observations. Funding for the TESS mission is provided by the NASA Explorer Program.

## DATA AVAILABILITY

The ground-based and TESS data used in this study are available upon request to the corresponding author.

## ORCID IDS

Atila Poro: 0000-0002-0196-9732  
 Farzaneh Ahangarani Farahani: 0000-0003-3740-7309  
 Esfandiar Jahangiri: 0000-0002-1576-798X  
 Ahmad Sarostad: 0000-0001-6485-8696  
 Mohammadjavad Gozarandi: 0000-0002-5131-6801  
 Marzie Haghgou: 0000-0002-1182-5871  
 Faezeh Abolhassani: 0000-0002-1235-4602  
 Amirhossein Fakhrabadi: 0000-0002-2735-7021  
 Anaël Wünsche: 0000-0002-6176-9847  
 Ramon Naves: 0000-0002-9463-9029  
 Pere Guerra: 0000-0002-4308-2339  
 Alessandro Marchini: 0000-0003-3779-6762  
 Mark Salisbury: 0009-0000-6809-7679  
 Roman Ehrenberger: 0000-0002-9720-9868

## REFERENCES

- [1] Mayor, M. & Queloz, D. 1995, Nature, 378, 355.
- [2] Gunther, S. & Guinan, E. F. 2021, AAS.
- [3] Fortney, J. J., Dawson, R. I., & Komacek, T. D. 2021, Journal of Geophysical Research (Planets), 126, e06629.
- [4] Charbonneau, D., Brown, T. M., Latham, D. W., et al. 2000, ApJ, 529, L45.
- [5] Hoyer, S., López-Morales, M., Rojo, P., et al. 2013, MNRAS, 434, 46.
- [6] Kane, S. R., Mahadevan, S., von Braun, K., et al. 2009, PASP, 121, 1386.
- [7] Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2014, Proc. SPIE, 9143, 914320.
- [8] Akeson, R. L., Chen, X., Ciardi, D., et al. 2013, PASP, 125, 989.
- [9] Poddaný, S., Brát, L., & Pejcha, O. 2010, New Astron., 15, 297.
- [10] Astropy Collaboration, Robitaille T. P., Tollerud E. J., et al., 2013, A&A, 558, A33.
- [11] Collins, K. A., Kielkopf, J. F., Stassun, K. G., et al. 2017, AJ, 153, 77.
- [12] Eastman, J., Gaudi, B. S., & Agol, E. 2013, PASP, 125, 83.
- [13] Salvatier, J., Wiecki, T. V., & Fonnesbeck, C. 2016, Astrophysics Source Code Library.
- [14] Noyes, R. W., Bakos, G. Á., Torres, G., et al. 2008, ApJ, 673, L79.
- [15] Hartman, J. D., Bakos, G. Á., Torres, G., et al. 2009, ApJ, 706, 785.

**Table 5.** Transit times of HAT-P-6b.

$T_c(BJD_{TDB})$	Error	Epoch	O-C	Ref./ETD Observer	$T_c(BJD_{TDB})$	Error	Epoch	O-C	Ref./ETD Observer
2454035.67648	0.00027	0	0.0000	[19]	2455160.75292	0.00034	292	0.0048	KPNO 2.1m
2454035.67652	0.00028	0	0.0000	[14]	2455430.45630	0.00180	362	-0.0008	[20]
2454347.76839	0.00042	81	0.0001	[19]	2455430.46118	0.00100	362	0.0041	R. Dřevěný
2454698.39160	0.00110	172	0.0017	[19]	2455430.46570	0.00130	362	0.0086	TRESCA
2454740.77668	0.00063	183	0.0039	UDEM	2455430.46701	0.00100	362	0.0099	P. Veres
2455064.42616	0.00119	267	0.0026	L. Brát	2456193.35362	0.00077	560	0.0055	Poddaný, Moudrá
2455064.42751	0.00089	267	0.0040	J. Trnka, M. Klos	2456208.76626	0.00063	564	0.0062	J. Garlitz
2455064.42843	0.00061	267	0.0049	Brát et al.	2457603.55511	0.00111	926	0.0145	J.L. Salto
2455064.43067	0.00134	267	0.0072	R. Dřevěný, T. Kalisch	2459495.37153	0.00076	1417	0.0153	M. Raetz

**Table 6.** Transit times of HAT-P-18b.

$T_c(BJD_{TDB})$	Error	Epoch	O-C	Ref./ETD Observer	$T_c(BJD_{TDB})$	Error	Epoch	O-C	Ref./ETD Observer
2454715.02174	0.00020	0	0.0000	[29]	2456455.56573	0.00075	316	0.0087	R. Naves
2455662.40408	0.00054	172	0.0024	V.P. Hentunen	2456780.53283	0.00089	375	0.0025	F.G. Horta
2455673.41967	0.00124	174	0.0019	[31]	2456780.53350	0.00073	375	0.0031	CAAT
2455695.45273	0.00081	178	0.0029	A. Ayiomamitis	2456780.53981	0.00128	375	0.0094	A. Carreno
2455706.46499	0.00077	180	-0.0009	C. Lopresti	2457116.52736	0.00073	436	0.0076	J. Gonzalez
2455706.46993	0.00080	180	0.0040	[31]	2457149.56917	0.00092	442	0.0013	M. Deldem
2455717.48233	0.00177	182	0.0004	A.L. Marrero	2457276.25646	0.00010	465	0.0040	[35]
2455728.49780	0.00100	184	-0.0002	V. Benishek	2457474.54607	0.00046	501	0.0048	Signoret, Pioppa
2455739.52081	0.00080	186	0.0068	F.C. Pecharrón	2457485.56256	0.00067	503	0.0053	Molina, Sureda
2456042.45973	0.00124	241	0.0044	F.G. Horta	2457485.56424	0.00084	503	0.0069	D. Molina
2456053.47276	0.00084	243	0.0014	[31]	2457507.59220	0.00019	507	0.0028	[32]
2456086.51856	0.00125	249	-0.0009	[31]	2457843.58653	0.00090	568	0.0077	Scaggiante, Zardin
2456097.53468	0.00074	251	-0.0008	T. Scarmato	2458559.62466	0.00111	698	0.0029	S. Dufoer
2456400.48452	0.00108	306	0.0077	V. Popov	2458730.37292	0.00129	729	0.0024	B. Andreas
2456411.49638	0.00084	308	0.0036	[31]	2459352.78320	0.00090	842	0.0061	[33]

**Table 7.** Transit times of HAT-P-22b.

$T_c(BJD_{TDB})$	Error	Epoch	O-C	Ref./ETD Observer	$T_c(BJD_{TDB})$	Error	Epoch	O-C	Ref./ETD Observer
2454930.22001	0.00025	0	0.0000	[17]	2457734.50182	0.00044	873	0.0137	V.P. Hentunen
2455601.57849	0.00079	209	0.0045	L. Brát	2457779.47248	0.00075	887	0.0133	D. Molina
2455614.43147	0.00126	213	0.0086	L. Brát	2457840.50512	0.00099	906	0.0138	M. Raetz
2455694.73397	0.00112	238	0.0056	S. Shadic, M. Butler	2458171.36382	0.00094	1009	0.0138	V.P. Hentunen
2455935.64851	0.00113	313	0.0036	S. Shadic	2458537.55631	0.00064	1123	0.0132	F. Campos
2455948.50056	0.00126	317	0.0068	G. Marino	2458550.40688	0.00053	1127	0.0149	J. Gaitan
2456298.63509	0.00098	426	0.0094	G.M. Schteinman	2458929.44872	0.00075	1245	0.0148	M. Theusner
2456327.53684	0.00066	435	0.0011	J. Lozano de Haro	2458945.50651	0.00071	1250	0.0115	M. Theusner
2456346.81517	0.00140	441	0.0061	[38]	2458945.51538	0.00117	1250	0.0204	A. Wunsche
2456356.45198	0.00059	444	0.0063	A. Ayiomamitis	2458945.51541	0.00055	1250	0.0204	Y. Jongen
2456391.78736	0.00088	455	0.0072	S. Shadic	2458945.51741	0.00106	1250	0.0224	M. Raetz
2456671.25040	0.00020	542	0.0071	[36]	2458974.42843	0.00083	1259	0.0234	Scaggiante, Zardin
2456687.31391	0.00055	547	0.0096	V.P. Hentunen	2459202.49043	0.00142	1330	0.0178	V. Perroud
2456706.58936	0.00090	553	0.0117	[37]	2459276.36752	0.00079	1353	0.0138	Scaggiante, Zardin
2456738.70941	0.00062	563	0.0095	[38]	2459276.37348	0.00044	1353	0.0198	M. Raetz
2457024.58745	0.00090	652	0.0000	CAAT	2459308.49480	0.00049	1363	0.0189	M. Bachschmidt
2457024.59630	0.00105	652	0.0089	R.N. Nogues	2459308.49839	0.00110	1363	0.0225	Y. Jongen
2457127.38972	0.00132	684	0.0112	A. Valdera					

**Table 8.** Transit times of HAT-P-12b.

$T_c(BJD_{TDB})$	Error	Epoch	O-C	Ref./ETD Observer	$T_c(BJD_{TDB})$	Error	Epoch	O-C	Ref./ETD Observer
2454187.85647	0.00061	-72	0.0012	[15]	2456732.59827	0.00063	720	-0.0003	[24]
2454216.77343	0.00023	-63	0.0006	[15]	2456735.81023	0.00029	721	-0.0014	[27]
2454419.19556	0.00020	0	0.0000	[15]	2456745.45299	0.00046	724	0.0021	G. Corfini
2454869.02462	0.00033	140	0.0007	[15]	2456745.45375	0.00074	724	0.0029	A. Valdera
2454897.94297	0.00057	149	0.0015	[15]	2456764.72820	0.00040	730	-0.0010	[27]
2454965.41502	0.00062	170	-0.0007	H. Kučáková	2456793.64833	0.00092	739	0.0016	[24]
2455347.76929	0.00021	289	-0.0006	[21]	2456806.49642	0.00062	743	-0.0026	[24]
2455630.51929	0.00011	377	0.0002	[24]	2456806.49815	0.00019	743	-0.0008	[24]
2455646.58184	0.00049	382	-0.0026	Š. Gajdoš, J. Világi	2456809.71238	0.00045	744	0.0003	[23]
2455649.79746	0.00020	383	0.0000	[27]	2457063.54386	0.00040	823	0.0001	M. Bretton
2455649.79770	0.00035	383	0.0002	[22]	2457124.59055	0.00039	842	-0.0014	M. Bretton
2455659.43649	0.00044	386	-0.0002	[24]	2457217.76896	0.00057	871	-0.0017	[27]
2455659.43650	0.00037	386	-0.0001	J. Ruiz	2457516.58394	0.00031	964	-0.0013	[25]
2455665.86227	0.00037	388	-0.0005	[22]	2457529.43692	0.00016	968	-0.0005	M. Bretton
2455669.07486	0.00080	389	-0.0010	[27]	2457831.46339	0.00037	1062	-0.0017	[25]
2455675.50436	0.00047	391	0.0024	[24]	2457834.67573	0.00042	1063	-0.0024	[25]
2455678.71382	0.00041	392	-0.0012	[27]	2457837.88977	0.00095	1064	-0.0014	[26]
2455678.71462	0.00047	392	-0.0004	[23]	2457847.53037	0.00025	1067	0.0000	M. Bretton
2455691.56661	0.00022	396	-0.0006	[24]	2457876.44558	0.00036	1076	-0.0023	[25]
2455694.78089	0.00029	397	0.0006	[22]	2457879.66160	0.00036	1077	0.0006	[25]
2455707.63244	0.00027	401	-0.0001	[24]	2457892.51113	0.00046	1081	-0.0021	[25]
2455720.48443	0.00015	405	-0.0003	[24]	2457892.51191	0.00012	1081	-0.0013	[25]
2455736.54999	0.00030	410	-0.0001	[24]	2457892.51258	0.00041	1081	-0.0006	[25]
2455964.67787	0.00059	481	0.0005	[24]	2457908.57706	0.00040	1086	-0.0014	[25]
2456006.44644	0.00062	494	-0.0007	V. Přibík	2457940.70759	0.00036	1096	-0.0015	[25]
2456006.44779	0.00012	494	0.0007	[24]	2458223.46181	0.00092	1184	0.0034	V. Hentunen
2456041.79018	0.00037	505	-0.0006	[23]	2458541.55104	0.00042	1283	-0.0002	Y. Jongen
2456051.42937	0.00048	508	-0.0006	[24]	2458570.46838	0.00042	1292	-0.0004	Y. Jongen
2456054.64202	0.00060	509	-0.0010	[24]	2458859.64077	0.00055	1382	-0.0034	V. Hentunen
2456070.70792	0.00035	514	-0.0004	[23]	2458904.62566	0.00033	1396	-0.0014	Y. Jongen
2456083.55936	0.00065	518	-0.0012	[24]	2458933.54260	0.00047	1405	-0.0020	M. Raetz
2456086.77329	0.00051	519	-0.0003	[23]	2458933.54500	0.00064	1405	0.0004	Y. Jongen
2456359.88215	0.00014	604	-0.0015	[28]	2458933.54535	0.00076	1405	0.0008	R.F. Auer
2456363.09540	0.00019	605	-0.0013	[28]	2458946.39354	0.00065	1409	-0.0033	M. Raetz
2456388.80064	0.00048	613	-0.0006	[24]	2458946.39547	0.00062	1409	-0.0013	Y. Jongen
2456388.80101	0.00081	613	-0.0002	[24]	2458946.39639	0.00043	1409	-0.0004	M. Bretton
2456462.70081	0.00050	636	-0.0008	[23]	2458946.39685	0.00104	1409	0.0000	F. Salvaggio
2456716.53165	0.00033	715	-0.0017	[24]	2458962.46093	0.00031	1414	-0.0012	M. Raetz
2456732.59741	0.00023	720	-0.0012	[24]	2459296.61925	0.00038	1518	-0.0011	Y. Jongen
2456732.59765	0.00011	720	-0.0010	[24]	2459309.47053	0.00046	1522	-0.0020	M. Bachschmidt
2456732.59796	0.00009	720	-0.0007	[24]	2459309.47209	0.00044	1522	-0.0005	Montigiani, Mannucci
2456732.59811	0.00028	720	-0.0005	[24]	2459354.45314	0.00034	1536	-0.0023	Y. Jongen

**Table 9.** Transit times of HAT-P-52b.

$T_c(BJD_{TDB})$	Error	Epoch	O-C	Ref./ETD Observer	$T_c(BJD_{TDB})$	Error	Epoch	O-C	Ref./ETD Observer
2455852.10326	0.00041	0	0.0000	[18]	2457727.30532	0.00049	681	0.0037	M. Bretton
2455995.29102	0.00031	52	0.0008	[47]	2457749.32972	0.00193	689	-0.0007	[48]
2457311.51042	0.00142	530	0.0017	[48]	2458060.48649	0.00155	802	-0.0002	[48]
2457336.29302	0.00277	539	0.0019	[48]	2458140.34268	0.00137	831	0.0017	[48]
2457336.29562	0.00164	539	0.0045	M. Bretton	2458140.34636	0.00214	831	0.0054	[48]
2457399.62421	0.00133	562	0.0004	[48]	2458429.47243	0.00066	936	0.0040	P. Guerra
2457413.39279	0.00091	567	0.0010	[48]	2458531.35748	0.00170	973	0.0060	M. Raetz
2457724.55292	0.00121	680	0.0049	R. Naves	2459167.44167	0.00098	1204	0.0097	S. Gudmundsson
2457727.30439	0.00107	681	0.0027	[48]	2459514.39070	0.00048	1330	0.0057	A. Wünsche

**Table 10.** Transit times of HAT-P-32b.

$T_c (BJD_{TDB})$	Error	Epoch	O-C	Ref./ETD Observer	$T_c (BJD_{TDB})$	Error	Epoch	O-C	Ref./ETD Observer
2454420.44637	0.00009	0	0.0000	[16]	2456600.55546	0.00017	1014	0.0010	[41]
2455796.45134	0.00092	640	-0.0001	Moudrá, Sobotka	2456600.55612	0.00033	1014	0.0016	P. Benni
2455839.45347	0.00101	660	0.0018	[41]	2456611.30423	0.00064	1019	-0.0003	F.G. Horta
2455843.75341	0.00019	662	0.0017	[41]	2456615.60935	0.00034	1021	0.0048	M. Perchak
2455843.75341	0.00019	662	0.0017	[21]	2456617.75617	0.00042	1022	0.0016	S. Shadick
2455845.90287	0.00024	663	0.0012	[41]	2456628.50585	0.00031	1027	0.0013	[41]
2455845.90287	0.00024	663	0.0012	[21]	2456637.10480	0.00023	1031	0.0002	[42]
2455845.90314	0.00040	663	0.0015	[41]	2456639.25687	0.00046	1032	0.0022	Aldi, Leo, Bozza
2455845.90314	0.00040	663	0.0015	[21]	2456656.45533	0.00045	1040	0.0006	[41]
2455852.35180	0.00073	666	0.0001	A. Brosio	2456656.45798	0.00064	1040	0.0033	R. Naves
2455856.65098	0.00029	668	-0.0007	S. Shadick	2456957.45714	0.00044	1180	0.0013	E.D. Alonso
2455858.80105	0.00039	669	-0.0007	S. Shadick	2456959.60773	0.00024	1181	0.0019	C.G.J. Dibon
2455860.95425	0.00034	670	0.0025	S. Shadick	2456972.50716	0.00069	1187	0.0013	F.G. Horta
2455867.40301	0.00073	673	0.0013	[41]	2456998.30682	0.00029	1199	0.0009	M. Salisbury
2455880.30170	0.00035	679	-0.0001	G. Corfini	2457009.05775	0.00080	1204	0.0017	[42]
2455880.30267	0.00033	679	0.0009	[41]	2457013.35882	0.00042	1206	0.0028	F. Campos
2455895.35249	0.00080	686	0.0006	[41]	2457024.10743	0.00046	1211	0.0014	[42]
2455895.35297	0.00016	686	0.0011	[41]	2457245.55959	0.00028	1314	0.0027	M. Bretton
2455897.50328	0.00033	687	0.0014	[41]	2457286.40911	0.00032	1333	0.0021	M. Salisbury
2455910.40274	0.00043	693	0.0008	[41]	2457303.60475	0.00055	1341	-0.0024	F. Signoret
2455912.54988	0.00147	694	-0.0020	R. Naves	2457342.30813	0.00054	1359	0.0009	M. Bretton
2455923.30295	0.00031	699	0.0010	[41]	2457372.41074	0.00069	1373	0.0034	D. Molina
2455940.50354	0.00109	707	0.0015	R. Naves	2457385.31188	0.00036	1379	0.0045	M. Bretton
2455942.65179	0.00113	708	-0.0002	[41]	2457396.05803	0.00091	1384	0.0006	[42]
2455942.65287	0.00064	708	0.0008	[41]	2457415.40279	0.00081	1393	-0.0047	J. Gaitan
2455942.65443	0.00051	708	0.0024	D. Zaharevitz	2457660.50761	0.00057	1507	-0.0008	R. Roy
2456155.50385	0.00026	807	0.0010	[41]	2457660.50931	0.00049	1507	0.0009	M. Deldem
2456157.65470	0.00072	808	0.0019	[41]	2457664.80650	0.00255	1509	-0.0019	M. Fowler
2456170.55416	0.00089	814	0.0013	F. Centenera	2457692.76393	0.00187	1522	0.0054	M. Fowler
2456177.00392	0.00025	817	0.0010	[40]	2457725.00958	0.00017	1537	0.0009	W. Kang
2456183.45361	0.00049	820	0.0007	[41]	2457759.41008	0.00034	1553	0.0013	V.P. Hentunen
2456183.45364	0.00085	820	0.0007	[41]	2457772.30878	0.00034	1559	-0.0001	M. Bretton
2456185.60375	0.00033	821	0.0008	[41]	2458004.51187	0.00057	1667	0.0022	F.G. Horta
2456211.40267	0.00214	833	-0.0004	[41]	2458021.71057	0.00027	1675	0.0008	O. Cooper
2456211.40361	0.00056	833	0.0006	[41]	2458088.36081	0.00032	1706	0.0008	P. Guerra
2456220.00440	0.00019	837	0.0013	[40]	2458107.71300	0.00071	1715	0.0029	[43]
2456226.45618	0.00102	840	0.0031	[41]	2458116.31178	0.00030	1719	0.0017	F. Bruno
2456228.60389	0.00057	841	0.0008	A. Carreño	2458367.86227	0.00074	1836	0.0012	B. Rodgers
2456232.90554	0.00028	843	0.0024	Kehusmaa, Harlington	2458376.46386	0.00042	1840	0.0028	G. Rousseau
2456237.20386	0.00029	845	0.0007	[42]	2458391.51025	0.00020	1847	-0.0009	M. Bretton
2456239.35339	0.00050	846	0.0002	J. Gaitan	2458404.40968	0.00081	1853	-0.0015	A. Wiünsche
2456239.35397	0.00052	846	0.0008	F.G. Horta	2458406.56109	0.00049	1854	-0.0001	V.P. Hentunen
2456243.65354	0.00086	848	0.0004	R. Naves	2458447.41287	0.00067	1873	0.0015	Y. Jongen
2456250.10439	0.00042	851	0.0012	[42]	2458490.41085	0.00040	1893	-0.0007	S.F. Dagot
2456254.40251	0.00063	853	-0.0007	R. Naves	2458503.31396	0.00031	1899	0.0024	Y. Jongen
2456254.40404	0.00022	853	0.0008	[41]	2458518.36601	0.00067	1906	0.0044	D. Laloum
2456265.15466	0.00048	858	0.0014	[42]	2458720.46171	0.00129	2000	-0.0007	V. Perroud
2456284.50460	0.00100	867	0.0013	[41]	2458739.81430	0.00382	2009	0.0019	[43]
2456297.39940	0.00069	873	-0.0040	R. Naves	2458821.51375	0.00044	2047	0.0010	Y. Jongen
2456514.55505	0.00065	974	0.0009	F.G. Horta	2458847.31493	0.00044	2059	0.0021	J. Dirscherl
2456527.45567	0.00030	980	0.0015	V.P. Hentunen	2458849.46424	0.00031	2060	0.0014	Y. Jongen
2456542.50522	0.00052	987	0.0010	[41]	2458862.36411	0.00035	2066	0.0012	Y. Jongen
2456542.50530	0.00018	987	0.0010	[41]	2459109.61612	0.00038	2181	0.0023	Y. Jongen
2456542.50538	0.00032	987	0.0011	[41]	2459120.36332	0.00054	2186	-0.0005	V.P. Hentunen
2456557.55814	0.00057	994	0.0038	Herrero, Pascual	2459178.41494	0.00085	2213	0.0009	F. Mortari
2456559.70548	0.00025	995	0.0011	P. Benni	2459197.76032	0.00152	2222	-0.0038	R. Dymock
2456572.60532	0.00018	1001	0.0009	[41]	2459221.41318	0.00066	2233	-0.0011	S. Gudmundsson
2456585.50553	0.00122	1007	0.0011	J. Mravík, J. Grnja	2459481.56450	0.00054	2354	-0.0007	Y. Jongen
2456598.40539	0.00017	1013	0.0009	[41]	2459535.31403	0.00113	2379	-0.0014	D. Daniel

**Table 11.** Transit times of HAT-P-33b.

$T_c(BJD_{TDB})$	Error	Epoch	O-C	Ref./ETD Observer	$T_c(BJD_{TDB})$	Error	Epoch	O-C	Ref./ETD Observer
2454429.93117	0.00055	-196	0.0021	[16]	2456723.08352	0.00047	464	0.0016	[44]
2454551.53949	0.00058	-161	0.0039	[16]	2456723.08417	0.00059	464	0.0023	[44]
2454794.75180	0.00037	-91	0.0030	[16]	2457039.26162	0.00046	555	0.0026	[44]
2454801.69988	0.00074	-89	0.0021	[16]	2457046.20981	0.00046	557	0.0018	[44]
2455110.92595	0.00022	0	0.0000	[16]	2457053.15793	0.00047	559	0.0010	[44]
2455607.77606	0.00060	143	0.0003	[16]	2457067.05749	0.00043	563	0.0027	[44]
2455614.72513	0.00036	145	0.0005	[16]	2457397.13183	0.00066	658	0.0020	[44]
2455857.93657	0.00067	215	-0.0013	S. Shadick	2458498.54660	0.00113	975	0.0085	P. Guerra
2455864.88110	0.00078	217	-0.0057	S. Shadick	2459172.58691	0.00093	1169	0.0009	V. Ferrando
2456024.71746	0.00120	263	0.0048	[46]	2459172.58950	0.00078	1169	0.0034	Y. Jongen
2456636.22181	0.00072	439	0.0018	[44]	2459172.59843	0.00148	1169	0.0124	A. Wünsche
2456716.13465	0.00069	462	0.0017	[44]	2459266.39915	0.00068	1196	0.0023	M. Raetz
2456716.13498	0.00065	462	0.0020	[44]					

- [16]Hartman, J. D., Bakos, G. Á., Torres, G., et al. 2011, ApJ, 742, 59.  
[17]Bakos, G. Á., Hartman, J., Torres, G., et al. 2011, ApJ, 742, 116.  
[18]Hartman, J. D., Bhatti, W., Bakos, G. Á., et al. 2015, AJ, 150, 168.  
[19]Szabo, G. M., Haja, O., Szatmary, K., et al. 2010, IBVS, 5919, 1.  
[20]Hébrard, G., Ehrenreich, D., Bouchy, F., et al. 2011, A&A, 527, L11.  
[21]Sada, P. V., Deming, D., Jennings, D. E., et al. 2012, PASP, 124, 212.  
[22]Lee, J. W., Youn, J.-H., Kim, S.-L., et al. 2012, AJ, 143, 95.  
[23]Sada, P. V. & Ramón-Fox, F. G. 2016, PASP, 128, 024402.  
[24]Mallonn, M., Nascimbeni, V., Weingrill, J., et al. 2015, A&A, 583, A138.  
[25]Alexoudi, X., Mallonn, M., von Essen, C., et al. 2018, A&A, 620, A142.  
[26]Yan, F., Espinoza, N., Molaverdikhani, K., et al. 2020, A&A, 642, A98.  
[27]Sariya, D. P., Jiang, I.-G., Su, L.-H., et al. 2021, Research in Astronomy and Astrophysics, 21, 097.  
[28]Baxter, C., Désert, J.-M., Tsai, S.-M., et al. 2021, A&A, 648, A127.  
[29]Hartman, J. D., Bakos, G. Á., Sato, B., et al. 2011, ApJ, 726, 52.  
[30]Esposito, M., Covino, E., Mancini, L., et al. 2014, A&A, 564, L13.  
[31]Seeliger, M., Kitze, M., Errmann, R., et al. 2015, MNRAS, 451, 4060.  
[32]Kirk, J., Wheatley, P. J., Louden, T., et al. 2017, MNRAS, 468, 3907.  
[33]Edwards, B., Ho, C., Osborne, H., et al. 2021, Astronomy: Theory, Observations and Methods, 2, 25.  
[34]Paragas, K., Vissapragada, S., Knutson, H. A., et al. 2021, ApJ, 909, L10.  
[35]Kokori, A., Tsiaras, A., Edwards, B., et al. 2022, ApJS, 258, 40.  
[36]Hinse, T. C., Han, W., Yoon, J.-N., et al. 2015, Journal of Astronomy and Space Sciences, 32, 21.  
[37]Baştürk, Ö., Hinse, T. C., Özavcı, İ., et al. 2015, Living Together: Planets, Host Stars and Binaries, 496, 370.  
[38]Turner, J. D., Pearson, K. A., Biddle, L. I., et al. 2016, MNRAS, 459, 789.  
[39]Mandel, K. & Agol, E. 2002, ApJ, 580, L171. doi:10.1086/345520  
[40]Gibson, N. P., Aigrain, S., Barstow, J. K., et al. 2013, MNRAS, 436, 2974.  
[41]Seeliger, M., Dimitrov, D., Kjurkchieva, D., et al. 2014, MNRAS, 441, 304.  
[42]Wang, Y.-H., Wang, S., Hinse, T. C., et al. 2019, AJ, 157, 82.  
[43]Zellem, R. T., Pearson, K. A., Blaser, E., et al. 2020, PASP, 132, 054401.  
[44]Wang, Y.-H., Wang, S., Liu, H.-G., et al. 2017, AJ, 154, 49.  
[45]Knutson H. A., Fulton B. J., Montet B. T., et al., 2014, ApJ, 785, 126.  
[46]Turner, J. D., Leiter, R. M., Biddle, L. I., et al. 2017, MNRAS, 472, 3871.  
[47]Wang, X.-Y., Wang, Y.-H., Wang, S., et al. 2021, ApJS, 255, 15.  
[48]Mallonn M., von Essen C., Herrero E., Alexoudi X., Granzer T., Sosa M., Strassmeier K. G., et al., 2019, A&A, 622, A81.

This paper has been typeset from a  $\text{\TeX}/\text{\LaTeX}$  file prepared by the author.