

TWIG BLIGHT OF JUNIPER

J. J. McRitchie and T. S. Schubert¹

At one time southern redcedar, *Juniperus silicicola* (Small) Bailey, was so plentiful that it occurred in dense forests on the west coast of Florida, particularly in the Cedar Key area. However, heavy lumbering for items such as wooden pencils virtually eliminated trees of commercial size. Southern redcedar is still valued for many landscaping uses, in part because of its cold hardiness and salt resistance. It is also popular as a living Christmas tree(1). Southern redcedar, and its very similar relative *J. virginiana* L. (eastern redcedar) are plagued by twig blight caused by the fungus *Phomopsis juniperovora* Hahn. Twig blight is found throughout the central and eastern United States wherever



Fig. 1. Segment of main stem of a southern redcedar seedling infected with *Phomopsis juniperovora*. Black pycnidia are visible on stem at top of segment, and pycnidia exuding cirrhi (spore horns or tendrils) are present on lower end of stem segment.

redcedar, common juniper, and many of their varieties are grown. The twig blight fungus has also been reported on arborvitae, cypress, Douglas fir, fir, hemlock, larch, redwood, white-cedar, and yew. Several juniper varieties are reported to be resistant to the fungus under field conditions (2).

SYMPTOMS. Tips of twigs affected by this disease first turn light to gray-green, then tan-brown, and eventually ashy gray. Tip blight may be followed by a progressive dying back of the entire branch and even girdling of the main stem at the axil of the infected branch. Small black fruiting structures (pycnidia) of the fungus may be observed on the dead tissue (Fig. 1).

Twig blight is primarily a disease of seedlings and nursery stock, although it is not unknown on larger trees. Usually, little damage occurs on trees over 5 years old. Junipers from cuttings of mature foliage generally have less of a problem with *Phomopsis* blight than seedlings or cuttings with juvenile foliage.

¹Plant Pathologists, Bureau of Plant Pathology, P. O. Box 1269, Gainesville, FL 32602.

DISEASE DEVELOPMENT. Spores from diseased tissue are splashed and/or wind-blown to healthy young needles. Infection may occur in a short period (7 hours) of 100 percent relative humidity. Infection is optimum at 24C, but disease development is favored by higher temperatures (32C). Symptoms may develop in as little as 3-5 days, but pycnidia are usually not seen for 3-4 weeks, after the tissue has dried. Fruiting bodies on dead tissue may continue to produce spores for as long as two years (4).

CONTROL. Seedling twig blight should be controlled in the nursery, before trees are planted out. Losses as high as 97 percent have been reported 3 years after planting out heavily diseased nursery seedlings (3). Twig blight in the nursery can be reduced by 1) spacing plants to provide good air movement and rapid drying; 2) avoiding planting in heavy shade; 3) avoiding handling wet plants; 4) pruning infected plants during dry weather; 5) planting resistant varieties (2); and 6) applying approved fungicides beginning with new shoot development and during periods of wet weather (2). Manzate 200, Zyban and benomyl are all EPA-registered for Phomopsis control on junipers. Disease-conducive conditions can occur year round in Florida.

SURVEY AND DETECTION. Look for the tips of young twigs and branches turning light green to brown, then ashy gray. Pycnidia of the fungus may be seen on the dry tissue.

LITERATURE CITED.

1. Bush, C. S. and J. F. Morton. 1969. Native trees and plants for Florida landscaping (revised edition). Florida Dept. Agric. Consumer Serv. Bull. 193:43-45.
2. Hartman, J. 1982. Juniper diseases. in Diseases of woody ornamental plants and their control in nurseries. (R. K. Jones and R. C. Lambe, eds.) N. C. State Univ., Raleigh. p60.
3. Hepting, G. H. 1971. Diseases of forest and shade trees of the United States. U.S.D.A. Forest Service Agr. Handbook 386:200-202.
4. Peterson, G. W. 1975. Phomopsis blight of junipers. in Forest nursery diseases in the United States. (G. W. Peterson and R. S. Smith, Jr., eds.). U.S.D.A. Forest Service Agr. Handbook 470:76-79.